

Things Need to Remember (Midterm 1)

$$\langle x|p\rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{\frac{ipx}{\hbar}} \quad ; \quad \langle p|x\rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{-\frac{ipx}{\hbar}}$$

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega}} (\hat{a}^\dagger + \hat{a})$$

$$\hat{p} = i\sqrt{\frac{m\omega\hbar}{2}} (\hat{a}^\dagger - \hat{a})$$

$$\hat{H} = \hbar\omega \left(\hat{N} + \frac{1}{2} \right) = \frac{\hat{p}^2}{2m} + V(\hat{r}) \quad , \quad (\hat{N} = \hat{a}^\dagger \hat{a})$$

$$\hat{a}^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle \quad ; \quad \hat{a} |n\rangle = \sqrt{n} |n-1\rangle$$


Schrödinger Picture : "state" depends on time, while operator not.

Heisenberg Picture : "Operator" depends on time, while state not.

Interacting Picture : "Both" operator & state depend on time.

Variational Method:
$$E_\lambda = \frac{\int dx \psi^*(x) \left[-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x) \right] \psi(x)}{\psi(x) \psi^*(x)}$$

WKB Method:
$$\psi(x) = \frac{A}{\sqrt{p(x)}} e^{\pm \frac{i}{\hbar} \int_{x_0}^x p(x') dx'} \quad , \quad p(x') = \sqrt{2m(E - V(x'))}$$



$$\begin{cases} \psi_{II}(x) \sim \frac{A}{\sqrt{p(x)}} e^{\frac{1}{\hbar} \int^x \sqrt{2m(V(x') - E)} dx} \\ \psi_I(x) \sim \frac{A}{\sqrt{p(x)}} e^{\frac{1}{\hbar} \int^x \sqrt{2m(V(x') - E)} dx} \\ \psi_{II}(x) \begin{cases} x \rightarrow x_1 : \frac{A}{\sqrt{p(x)}} \cos\left(\frac{1}{\hbar} \int_{x_1}^x p(x') dx - \frac{\pi}{4}\right) \\ x \rightarrow x_2 : \frac{A}{\sqrt{p(x)}} \cos\left(\frac{1}{\hbar} \int_{x_2}^x p(x') dx + \frac{\pi}{4}\right) \end{cases} \end{cases}$$

⇒ Quantization condition:

$$\int_{x_1}^{x_2} \sqrt{2m(E - V(x'))} dx' = \left(n + \frac{1}{2}\right) \hbar \pi$$